

Jet Structure Topical Update

Rosi Reed and Dennis Perepelitsa

Jet structure in the RHIC and LHC era

- Jet measurements in heavy-ion collisions have evolved in the last year
 - Truly the era of jet structure
 - HEP inspired observables
- Which observables should the next generation jet (and Y!) detector focus on?
 - \square $\sqrt{S_{NN}}$ dependence
 - 200 GeV vs 2.76, 5 TeV
 - Focus on differential measures at 200 GeV
- Jet grooming, jet fragmentation, jet mass, N_{subjets} , γ-jet, jet shape, jet girth





I-PREL-101580

sPHENIX Jet structure goals

- Tracking design is now realized
 - Key aspect for looking at intra-jet correlations
 - Jet selection from calorimeters which removes autocorrelations
- Clustering needs to be continued to be evaluated
 - \blacksquare Calorimeter clustering \rightarrow Important for γ observables
 - Requires track-cluster matching evaluation is importing
- Keep up with the changing face of jets in heavy-ion physics
 - \square Some observables still key (γ-jet)
 - Some observables are new (jet grooming)
 - Which are the best for day one physics analyses (most differential, cleanest comparison LHC results)

sPHENIX Jet structure goals

- Evaluate jet/single particle performance
 - Systematic studies versus jet kinematics and jet-finding characteristics are needed
 - Benchmark with new techniques given person-power/interest, for example evaluating particle flow with finalized tracking design
- Determine the best way to interface with JETSCAPE
 - http://jetscape.wayne.edu/jetscape/
 - Theory input will be valuable in determining the next generation of jet observables
- Jet Structure meeting next week
 - **I** Follow up on tracking simulation, γ clustering, new observables

Example Observable z_g

- Observable: Momentum fraction carried by the subleading branch of first splitting
 - Independent of flavor (q/g fraction irrelevant)
 - Tracking only observable → Possible to evaluate now!

